



Radiation Research Trust Conference
September 2008

Mobile Phones and Health – 15 Years of Research

EMBARGOED UNTIL MONDAY 8TH SEPTEMBER 2008

Radiation Research Trust Conference

September 2008

Mobile Phones and Health – 15 Years of Research

a report by Graham Philips of Powerwatch

on behalf of the Radiation Research Trust

Background	2
Mobile / cordless phones and brain cancer.....	2
<i>Lennart Hardell and Swedish research</i>	2
<i>Flaws, recall bias and selection bias</i>	3
<i>The Interphone Project</i>	4
<i>Other research</i>	6
Mobile phones and other tumours.....	6
<i>Parotid / salivary gland tumours</i>	6
<i>Other Tumours</i>	6
Mobile phone radiation and fertility	7
Mobile phone radiation and EEG or neurological effects.....	7
<i>Neurological effects</i>	7
<i>EEG and rCBF effects</i>	7
<i>Other effects</i>	8
Mobile phone radiation and DNA strand breaks / cell damage.....	8
<i>DNA strand breaks and cell mutation / death</i>	8
<i>Heat shock protein type effects</i>	9
<i>Nerve damage and signalling effects</i>	9
References.....	9

Background

Mobile phones have revolutionised communications throughout the world. Texting has resulted in a new language barely understood by the pre-mobile phone generation. It is hard to imagine the modern business and younger social world without the mobile phone. However, research from across the globe is showing that long-term usage of mobile phones can be damaging to human health. It is associated with brain tumours, memory and concentration problems, and experiments on rats have found signs of early onset dementia (similar to Alzheimer's - [Salford 2003](#)). The UK Department of Health has also issued adviceⁱ stating that people should attempt to keep their calls as short as possible and that children and young people under the age of 16 should only use their phones in emergencies, based on the current uncertainties in the existing scientific literature.

This paper seeks to provide an overview of the research undertaken over the last fifteen years assessing the health effects of mobile phones. A full list of papers referenced in this document is included below.

Mobile / cordless phones and brain cancer

There has been repeated media coverage in the last few years that mobile phones can increase the risks of both malignant and benign forms of brain tumours. There has been vehement criticism of this coverage, claiming that there is no scientific evidence to support such an association and that reporting a link is inappropriate at best, and irresponsible scaremongering at worst.

In actual fact, there is now a large amount of epidemiological literature assessing the risk of mobile phone usage and brain cancer. In reality we would not expect studies to have found a link at this point in time, as the latency period (time between exposure to cause and diagnosis of cancer) of most brain tumours is between 15 and 25 years, and mobile phones have only been in widespread usage for about 10 years. Despite this, a number of papers are now showing signs of a significantly increased risk of brain cancer incidence from long term usage (over 10 years).

Lennart Hardell and Swedish research

Oncologist Lennart Hardell and his colleagues have been researching this association for about 10 years and have published numerous papers covering their findings. Their preliminary work at the turn of the century found a statistically significant 1.4-fold increase in risk for brain tumours (not sub-categorised) on the same side of the head as the mobile phone was used [[Hardell 1999](#), [Hardell 2000](#)].

In 2002, looking at 1617 patients histopathologically diagnosed with brain tumours, they found that use of analogue mobile phones was associated with a significant 30% increase in risk for brain tumours (overall). This increased to 80% when only looking at patients who had used their phone for over 10 years, and further to 150% (2.5 times as likely to develop a brain tumour) when side of the head was taken into account. For acoustic neuromas, the increase in risk was 250% [[Hardell 2002](#)]. This was followed in 2003 by a separate analysis that found that Astrocytomas also had a significantly increased risk of 80% [[Hardell Feb 2003](#)], and a subsequent paper finding a 3.5-fold risk of Acoustic Neuroma from mobile phone use (CI 1.77-6.76) [[Hardell Mar 2003](#)].

By 2005 most papers generally sub-categorised brain tumour risk from mobile phone usage into acoustic neuromas, meningiomas and gliomas. Hardell produced a paper in 2005 analysing the increase in risk for acoustic neuromas and meningiomas. With a very good response rate (85%) he found that, for a mobile phone usage of greater than 10 years, the odds ratio for meningiomas

was 2.1 (CI 1.1-4.3) and for acoustic neuromas this was split further into digital mobile phones (2.0 - CI 1.05-3.8) and analogue mobile phones (4.2 - CI 1.8-10) [Hardell 2005]. By this stage his results were consistently pointing towards a possible doubling in risk for mobile phone users who had not used their phone for as long as the typical latency period for the tumours. Hardell also published one of the only papers to date looking at risk of non-Hodgkin's lymphoma, finding a 6-fold increase for over 10 years of use. He did highlight however that there were very few cases and that the results should be interpreted with a fair degree of caution [Hardell Sept 2005].

In February 2006 Hardell published a paper using more recent diagnoses (patients diagnosed between 2000 and 2003), and found that the increase in risk was steadily strengthening in magnitude and statistical significance as the length of phone usage was increasing. For malignant tumours he found that the OR for analogue phone use was 2.6 (CI 1.5-4.3), for digital phone use was 1.9 (CI 1.3-2.7), and for cordless phone use was 2.1 (CI 1.4-3.0). Looking at patients who had used their phones for 10 years or more this increased to 3.5 (CI 2.0-6.4), 3.6 (CI 1.7-7.5) and 2.9 (CI 1.6-5.2) respectively [Hardell Feb 2006]. This work was followed up in October 2006 looking specifically at acoustic neuromas (a benign form of brain tumour), astrocytomas and non-Hodgkin's lymphomas. He found acoustic neuromas and the higher grades of astrocytoma (Grades III and IV) to have significant increases from all forms of mobile and cordless phone usages (around 50% increase in risk in each case), which increased further for those who had used their phone greater than 10 years. However, he found no increase for lower grade astrocytomas (Grade I and II), and no increase for non-Hodgkin's lymphomas in contrast to his paper from the previous year [Hardell Oct 2006].

Hardell published a meta-analysis in September 2007 of the existing literature to date (2 cohort studies and 16 case-control studies). His findings were a 140% increase in risk for benign acoustic neuromas (CI 1.1-5.3) and a 100% increase in risk for malignant gliomas (CI 1.2-3.4), with further increased risks when looking at ipsilateral exposure (exposure on the same side of the head as the tumour) [Hardell Sept 2007]. His summary so far on all the work (including work from other researchers) is that "Results from present studies on use of mobile phones for > or =10 years give a consistent pattern of increased risk for acoustic neuroma and glioma. The risk is highest for ipsilateral exposure.

Flaws, recall bias and selection bias

As Hardell is one of the only researchers consistently finding an effect, he has received a lot of criticism for the suspected quality (or lack of) his work. The primary criticism that has been directed at his work is that of recall bias (failure of the mobile phone users to correctly remember the amount of usage and the side of the head). However (and primarily for the Interphone study), this has been addressed by Vrijheid in a study that found that heavy users generally overestimated their use and light users generally underestimated their use [Vrijheid 2006]. As the high risk category is the heavy users, an overestimation of use would imply the risk is actually attributed to lower usage, which will lead to an underestimation of risk. Vrijheid also found a small element of inaccuracy of users recalling which side of the head they used their phone which may contribute to a slight overestimation of risk in the highest user category [Vrijheid 2008]. The other Interphone study to look at recall bias found that users were more accurate at recalling phone usage in terms of number of calls made rather than total phone usage, but without comment as to how this was likely to effect the reported risk [Samkange-Zeeb 2004]. One of the Interphone papers looked at selection bias, and found that approximately 10% less cases responded than controls (70% and 80% response rates respectively) [Lahkola 2005]. If this difference is real, it would lead to an overestimation in risk, but it is only based on one of the Interphone studies. Lloyd Morgan's pooled analysis found only 60% of controls responded on average (see the section on flaws of the Interphone Project below) - if so, this would instead lead to an underestimation in risk.

The Interphone project

The Interphone project, initiated in 1999, is a multi-national initiative coordinated by the International Agency for Cancer Research (IACR). Its aim is to fully examine the association between long term mobile phone usage and increases in risk for all forms of brain cancer. The results of the project were expected to be published in 2005, but the findings were conflicting and the researchers involved in the project were divided into three opinions on how the results should be summarised: a) that there was a possible increase in brain tumour risk that is real and warrants further investigation, b) that the increase is an artefact due to bias in the study, and c) that with the data currently available it is not possible to tell whether a) or b) are correct, with both being a feasible possibility. Louis Slesin has covered the various updates in the Interphone project in great detail on his Microwave News website. The results of the Interphone project have still (as of August 2008) not been published.

Unfortunately, there are a number of flaws with the design of the studies in the Interphone project. One of the major flaws is a lack of inclusion in most of the studies of any form of recognition of digital cordless phone users. With the handset similar to a phone and the base unit exposing users all the time whilst they are in the building, this becomes a very large confounder for using "length of phone usage" as a metric of exposure level.

Lloyd Morgan, an American researcher and electronic engineer (and currently a member of the board of directors at the Central Brain Tumour Registry of the United States), has been writing about the Interphone papers as each one is published, and has explained the major flaws with the Interphone protocol in great detail. The summary of the 6 primary flaws are listed below:

Flaw 1: Selection bias

The first flaw is called selection bias. This is likely to be the result of the low percentage of controls that participated in the studies (weighted average of 59%). Think about being randomly selected for a cellphone study. You are told you will be asked to answer a long questionnaire. If you use a cellphone you are more likely to agree to participate than if you do not use a cellphone. If this happens it is called selection bias. Selection bias will result in an underestimation of risk.

Flaw 2: Inclusion of tumours outside the cellphone's radiation plume

The second flaw is the inclusion of all brain tumours without regard to their location. Because the cellphone's radiation plume only penetrates a short distance into the head, nearly all of this radiation is absorbed by the temporal lobe, the acoustic nerve, or the parotid gland. Even when cellphone exposure of one side of the head is considered on the side where the cellphone was held, a substantial portion of half the brain is unexposed (the opposite side is completely unexposed). Studies that include brain tumours outside of the cellphone radiation plume contribute to an underestimation of the risk of brain tumours.

Flaw 3: Latency time and definition of regular user

The third flaw is the definition of "regular" cellphone use in relation to a reasonable latency time. "Regular" cellphone use is defined as use of a cellphone on average once per week for at least 6 months. Exposure within 1 year of the diagnosis date is not considered. The result of this definition, combined with the incredibly fast rate of new cellphone users, is to overweight "regular" users with an incredibly large group of short-term users - far too short a time to expect a tumour to be diagnosed.

The latency time for brain tumours is between 15 and 25 years. For the Interphone studies, using weighted averages for cases or controls, we see that 6.3% of cases and 6.4% of controls have used a cellphone for 10 years or more, and 18% of cases and 21% of controls have used a cellphone for 5 years or more. (Weighted average of 13 Interphone brain tumour studies - 10 Interphone studies are included and 3 are excluded because they are pooled studies which overlap the other 10). For a reasonable latency time, it would be unlikely to find any risk of tumours, given the percentage of cases and controls. Yet some of the Interphone studies are already finding a risk. Because such a large percentage of "regular" users have used a cellphone for an unreasonably short latency time the reported results for < 10 years as well as for > 10 years (6.3% of cases) are an underestimation of risk.

Flaw 4: Children and young adults are not included in interphone studies

The Interphone Protocol states that cases be between 30 and 59 years of age. While a few studies have included cases as young as 20, the non-inclusion of < 20 year olds is likely to result in an underestimation of risk. Research has shown that children's brains (due to skull formation through the childhood years) absorb a greater proportion of the radiation emitted by a mobile phone [Ghandi 1996, Ghandi 2002, Christ 2005, de Salles 2006, Wiart 2008]. They are also likely to be at a greater risk due to their higher rate of cell division (than adults). It is generally accepted that teenagers and young adults are the primary users of mobile phones.

Flaw 5: Cellphone's radiated power

It is reasonable to expect that risk of a tumour from a cellphone, after a reasonable latency time, would be the cellphone's power multiplied by cumulative time of use. In the early days of cellphone use all cellphones used analogue technology. These always radiated a fixed amount of power (~2 Watts). Analogue cellphone use has been totally displaced by digital cellphones. Digital cellphones have a feature called Automatic Power Control or APC. At the beginning of a call the cellphone radiates maximum power (~2 Watts) but quickly reduces the power so the radiated power is sufficient to have a reliable link to the cell tower (AKA masts or base stations). The result is that cellphones radiate far less power in urban areas compared to rural areas. This is because cell phone towers are much closer in urban areas compared to rural areas so the cellphone radiates less power in urban areas and more power in rural areas. When rural and urban cellphones are not reported separately the result is an underestimation of risk.

Flaw 6: Number of cases included in a study

The weighted average time in these 10 studies for a case to be eligible for inclusion in the study was only 2.6 years. When one considers 4 of the 5 previous flaws, it becomes obvious that such a short period of time for eligibility will result in too few cases to resolve these flaws. For example, if tumours were limited only to the exposed region of the brain then there would be far fewer cases; if a reasonably long latency time was included, again there would be far fewer cases; if children had been included there would have been more cases; and, if rural users were to be compared to the far larger number of urban users, a much larger number of cases would need to be eligible to participate in the Interphone Study.

A number of the Interphone studies have found statistically significant *protective* effectsⁱⁱⁱ from mobile phone usage [Christensen 2005, Lonn 2005, Schuz Mar 2006, Schuz Dec 2006, Klaeboe 2007, Lahkola 2007, Lahkola 2008], and despite these flaws some have still found statistically significant increase in risk for the heavier group of mobile phone users [Lonn Nov 2004, Schoemaker 2005, Hepworth 2006, Hours 2007, Takebayashi 2008].

A number of the papers have not shown this unlikely looking protective effect and have still not shown an increase in risk [Christensen 2004, Takebayashi 2006], but it is important to remember that with

around 5% of the cases having used their phone for 10 years or more we would not expect to see a risk anyway (as the latency period for these tumours tends to be between 15 and 25 years). The fact that any papers are showing a risk is very concerning.

Other research

There have been a number of other issues on an individual study basis, such as the highly publicised Danish cohort study towards the end of 2006 [[Schuz_Mar_2006](#), [Schuz_Dec_2006](#)] which was hyped as being a "definitive study", containing some 400,000 people in the dataset. Sadly, the classification of these subjects was extremely misleading: Firstly, only contract users were considered as those were the only users that it was possible to identify. All "pay as you go" users were classified as "non-mobile phone users" and effectively moved into the control group - this would lead to an underestimation of risk. Originally the authors had 720,000 records, of which 100,000 were removed (quite validly) due to duplication. Out of the remaining 620,000, a further 200,000 were removed because they "couldn't identify the users" as the contracts were corporate and not linked to any specific individual. By the authors' own admission these were likely to be the heaviest phone users in the dataset, so another 33% of the heaviest users they were looking for were moved in to the "non-user" control group, which would also lead to an underestimation of the risk. Strangely, the study's findings then highlighted a significant protective effect (7 of the 18 data points had a statistically significant reduction in risk), which disappeared in the highest usage category. One reason for this may be the flaws detailed by Lloyd Morgan above for the Interphone project papers that would expect a significantly reduced odds ratio (OR)ⁱⁱⁱ in cases when compared to controls. If these are controlled for in the statistical calculations, we end up finding an increase in risk of 30% for the heaviest users, reaching borderline significance.

There have been a number of other epidemiological papers published over the last 10 years looking at mobile phones and brain cancer, but most of these have failed to find an effect. This is unsurprising, as the studies were published a number of years ago and the cases involved had used their phones for less than 5 years - far too short a period to find any risk of brain tumours [[Muscat_2000](#), [Johansen_2001](#), [Inskip_2001](#), [Muscat_2002](#), [Auvinen_2002](#), [Cook_2003](#), [Lonn_Jan_2004](#)]. Despite this, two of the studies found increases for brain cancer that were of borderline significance [[Muscat_2000](#), [Auvinen_2002](#)]. More recently Kan performed a meta-analysis on 9 case-control studies, finding no increase of risk overall but a statistically significant increase in risk (OR 1.25) for those who have used their phone for more than 10 years (CI 1.01-1.54)^{iv} [[Kan_2008](#)].

Mobile phones and other tumours

Parotid / salivary gland tumours

An Israeli study this year found a statistically significant association between mobile usage and parotid gland (salivary gland) tumours [[Sadetzki_2008](#)]. Whilst the findings show a 60% increase in risk, other research from the previous 5 years failed to find an increase [[Auvinen_2002](#), [Hardell_2004](#), [Lonn_2006](#)]. That said, this is another tumour with a long typical latency period (20 years), so any effect would be unexpected and should be treated with concern at this stage.

Other Tumours

Other tumours that have been assessed but found to have no association with mobile phone usage are central nervous system tumours and arrhythmia-related heart diseases [[Johansen_2004](#)], and malignant melanomas of the eye [[Johansen_2002](#)]. However, Stang published a paper the previous year

finding a statistically significant and considerable increase in risk for uveal melanoma (a malignant form of eye cancer - OR 4.2 - CI 1.2-14.5) [Stang 2001].

Mobile phone radiation and fertility

The other issue to have received a large amount of media attention recently is a possible link between mobile phone usage and male infertility. Five key papers have found statistically significant effects on fertility from heavy mobile phone usage [Baste 2008, Agarwal 2008, Eroglu 2006, Fejes 2005, Grajewski 2000]. Lennart Hardell published a paper in 2007 looking at phone use and testicular cancer, where an association was not found [Hardell April 2007].

Mobile phone radiation and EEG or neurological effects

Mobile phone usage has expanded enormously over the last 10 years, and teenagers / young adults are now amongst the heaviest users. There have been reports that concentration, behaviour, memory and other cognitive performance have been negatively affected by mobile phones and their base stations. If these are true, extensive mobile phone use could be responsible for behavioural and performance issues in children of school age. A study published in July 2008 found that children exposed prenatally and early postnatally to mobile phones had an 80% increase in behavioural problems (CI 1.45-2.23) [Divan 2008]. Other research into prenatal exposure has found that mobile phone exposure has a significant effect on foetal heart rate and cardiac output [Rezk 2008]. The following is a brief summary of the research looking into neurological effects from mobile phone or other pulsed radio frequency (RF) electromagnetic field exposure.

Neurological effects

With two exceptions (a Chinese study that found reaction times decreased from pulsed RF exposure [Cao 2000], and a recent study from Germany finding no effect on cognitive function or sleep [Fritzer 2007]), research has found that exposure to mobile phone or other pulsed RF radiation exposure improved reaction times [Preece 2005, Koivisto Feb 2000], general cognitive performance [Edelstyn 2002], and high intensity memory tasks [Koivisto June 2000, Krause Dec 2000, Krause March 2006]. A Turkish paper from 2006 also found significant temporary hearing loss from mobile phone exposure [Oktay 2006].

The precise implications and cause (if the association is truly causal) is not clear, but it is clear that the pulsed radiofrequency electromagnetic fields produced by mobile phones is having some effect on attention, concentration, memory, reaction times and general cognitive performance.

EEG and rCBF effects

Sleep, EEG (electroencephalograph)^v and waking rCBF^{vi} (regional cerebral blood flow) have been studied in relation to RF exposure for a decade now, and the majority of papers published have found some form of effect. Whilst a Finnish study failed to find any effect on sleep or other cognitive function from pulsed RF exposure [Haarala 2007], most other papers have found significant effects on sleep [Hung 2007, Huber 2005, Huber 2002, Huber 2000, Borbely 1999, Andrzejak 2008]. Interestingly, two of these papers found the effect was only present when the exposure was pulsed (amplitude modulated) [Huber 2005, Huber 2002]. One early paper actually found that sleep quality (measured by the amount of participants' broken sleep) actually improved [Borbely 1999].

Whilst some papers were inclusive or inconsistent [Krause 2007, Papageorgiou 2006], a number of studies have now demonstrated reversible EEG and rCBF alterations from exposure to pulsed RF exposure [Aalto 2006, Krause June 2006, D'Costa 2003, Kramarenko 2003]. German research from 2006 found that

statistically significant EEG changes could be consistently found, but only in a relatively low proportion of study participants (12 - 30%) [[Bachman 2006](#)].

Other effects

Mobile phones have been negatively associated with subjectively reported symptoms such as headaches, nausea, concentration issues, dizziness, blurred vision and a number of other symptoms that tend to be grouped under the umbrella category Electromagnetic Hypersensitivity. A number of double blind^{vii} provocation studies have failed to find the association, though there are a number of theorised flaws on the methods by which traditional provocation studies are carried out.

Bruce Hocking published 3 studies at the beginning of this decade looking specifically at finding objective signs on people subjectively suffering from some form of general dysaesthesiae (impairment of any of the senses), finding significant signs of nerve damage in those suffering [[Hocking 2001](#), [Hocking 2002](#), [Hocking 2003](#)]. These findings support the possibility, and provide a plausible mechanism, of subjective symptoms that some mobile phone users report to suffer from. The papers are all from only one author, and there are no papers that have attempted to replicate these findings, which now seem long overdue.

Mobile phone radiation and DNA strand breaks / cell damage

A superb 291-page summary into existing work by a number of laboratories across Europe summarised most of the existing cellular literature on Radio Frequency (RF) and extremely low frequency electromagnetic fields (ELF EMFs). They also conducted a considerable amount of extra research, publishing their results as a 291-page report under the European Union programme "Quality of Life and Management of Living Resources" as the REFLEX report [[REFLEX report 2004](#)].

DNA strand breaks and cell mutation / death

As early as 1996 researchers have found that radiofrequency EMFs are capable of causing cellular damage, when scientists from Canada found that prolonged (24 hour) exposure to 900 MHz radiation damaged the lungs of live pigs [[Singh 1996](#)]. Since then, a number of *in vivo* experiments have found mobile phone or simulated mobile phone radiation exposure can cause cell damage, reactive oxygen species formation (which are the primary cause of DNA strand breaks), and cell death [[Oktem 2005](#), [Oral 2006](#), [Ferreira 2006](#), [Panagopoulos 2007](#), [Yan 2008](#)].

It is often claimed that there is insufficient energy in microwave frequency radiation to cause DNA strand breaks (as in the case of, for example, ionising radiation). What is less known is that the primary cause of DNA strand breaks is not high energy radiation, but a by-product of ROS (reactive oxygen species) creation. A number of *in vitro* experiments have found an association between RF exposure and ROS production, and then subsequent DNA single and double strand breaks [[Nikolova 2005](#), [Friedman 2007](#), [Yao May 2008\(1\)](#), [Yao May 2008\(2\)](#)], though some papers have failed to replicate this effect [[Lantow 2006](#), [Valbonesi 2008](#)].

There have also been a number of other cellular effects found, from cell mutations (such as micronuclei formation and cellular aneuploidy) [[D'Ambrosio 2002](#), [Hoyto 2007](#), [Mazor 2008](#), [Schwarz 2008](#)], and even impaired cell repair or cell death [[Ioubert 2008](#), [Manti 2008](#)]. A study published by Belyaev in 2006 found that whilst he could not induce DNA double strand breaks, exposure to 900 MHz GSM radiation induced changes in gene expression in rat brain cells [[Belyaev 2006](#)].

Heat shock protein type effects

Heat shock proteins (also known as "stress proteins") are known for increasing their expression^{viii} when their cells are exposed to elevated stress. It is known that microwave frequency radiation sufficient to heat tissue triggers an increase in heat shock protein response. Research has found effects in both sub-thermal levels of RF exposure, and levels of response beyond those expected from just heating effects for higher levels of exposure [Velizarov 1999, Leszczynski 2002, de Pomerai 2003, Czyz 2004, Belyaev 2005, Nylund 2006, George 2008].

Nerve damage and signalling effects

Finally, a number of papers have found effects on nerve damage, calcium pathways, and neurotransmitter expression from exposure to pulsed microwave radiation [Lai 1989, Lai 1994, Donnellan 1997, Leszczynski 2002, Salford 2003, Wang 2004, Wang Mar 2005, Wang Sept 2005, Rao 2008]

Conclusion

This report highlights the need for precaution based on the overwhelming evidence that exposure to mobile phone radiation is able to cause adverse health effects in a high proportion of the general population. With a possible increased risk of brain cancers, reduced fertility, concentration issues and neurological damage, the consequences of inaction could be both serious and wide-reaching, affecting a large proportion of the general population.

References

Aalto S et al, (July 2006) *Mobile phone affects cerebral blood flow in humans*, J Cereb Blood Flow Metab. 2006 Jul;26(7):885-90

Agarwal A et al, (January 2008) *Effect of cell phone usage on semen analysis in men attending infertility clinic*, Fertil Steril. 2008 Jan;89(1):124-8

Andrzejak R et al, (August 2008) *The influence of the call with a mobile phone on heart rate variability parameters in healthy volunteers*, Ind Health. 2008 Aug;46(4):409-17

Auvinen A et al, (May 2002) *Brain tumors and salivary gland cancers among cellular telephone users*, Epidemiology. 2002 May;13(3):356-9

Bachmann M et al, (2006) *Integration of differences in EEG Analysis Reveals Changes in Human EEG Caused by Microwave*, Conf Proc IEEE Eng Med Biol Soc. 2006;1:1597-600

Baste V et al, (April 2008) *Radiofrequency electromagnetic fields; male infertility and sex ratio of offspring*, Eur J Epidemiol. 2008 Apr 16 [Epub ahead of print]

Belyaev IY et al, (April 2005) *915 MHz microwaves and 50 Hz magnetic field affect chromatin conformation and 53BP1 foci in human lymphocytes from hypersensitive and healthy persons*, Bioelectromagnetics. 2005 Apr;26(3):173-84

Belyaev I et al, (May 2006) *Exposure of rat brain to 915 MHz GSM microwaves induces changes in gene expression but not double stranded DNA breaks or effects on chromatin conformation*, Bioelectromagnetics. 2006 May;27(4):295-306

- Borbely AA et al**, (November 1999) *Pulsed high-frequency electromagnetic field affects human sleep and sleep electroencephalogram*, *Neurosci Lett*. 1999 Nov 19;275(3):207-10
- Cao Z et al**, (March 2000) *Effects of electromagnetic radiation from handsets of cellular telephone on neurobehavioral function*, *Wei Sheng Yan Jiu*. 2000 Mar 30;29(2):102-3
- Christ A, Kuster N**, (2005) *Differences in RF energy absorption in the heads of adults and children*, *Bioelectromagnetics*. 2005;Suppl 7:S31-44
- Christensen HC et al**, (February 2004) *Cellular telephone use and risk of acoustic neuroma*, *Am J Epidemiol*. 2004 Feb 1;159(3):277-83
- Christensen HC et al**, (April 2005) *Cellular telephones and risk for brain tumors: a population-based, incident case-control study*, *Neurology*. 2005 Apr 12;64(7):1189-95
- Cook A et al**, (June 2003) *Cellular telephone use and time trends for brain, head and neck tumours*, *N Z Med J*. 2003 Jun 6;116(1175):U457
- Czyz J et al**, (May 2004) *High frequency electromagnetic fields (GSM signals) affect gene expression levels in tumor suppressor p53-deficient embryonic stem cells*, *Bioelectromagnetics*. 2004 May;25(4):296-307
- D'Ambrosio G et al**, (January 2002) *Cytogenetic damage in human lymphocytes following GSMK phase modulated microwave exposure*, *Bioelectromagnetics*. 2002 Jan;23(1):7-13
- D'Costa H et al**, (December 2003) *Human brain wave activity during exposure to radiofrequency field emissions from mobile phones*, *Australas Phys Eng Sci Med*. 2003 Dec;26(4):162-7
- de Pomerai DI et al**, (May 2003) *Microwave radiation can alter protein conformation without bulk heating*, *FEBS Lett*. 2003 May 22;543(1-3):93-7
- de Salles AA et al**, (2006) *Electromagnetic absorption in the head of adults and children due to mobile phone operation close to the head*, *Electromagn Biol Med*. 2006;25(4):349-60
- Divan H et al**, (May 2008) *Prenatal and Postnatal Exposure to Cell Phone Use*, *Epidemiology*. 2008 May 7 [Epub ahead of print]
- Donnellan M et al**, (July 1997) *Effects of exposure to electromagnetic radiation at 835 MHz on growth, morphology and secretory characteristics of a mast cell analogue, RBL-2H3*, *Cell Biol Int*. 1997 Jul;21(7):427-39
- Edelstyn N, Oldershaw A**, (January 2002) *The acute effects of exposure to the electromagnetic field emitted by mobile phones on human attention*, *Neuroreport*. 2002 Jan 21;13(1):119-21
- Erogul O et al**, (October 2006) *Effects of electromagnetic radiation from a cellular phone on human sperm motility: an in vitro study*, *Arch Med Res* 37(7):840-3
- Fejes I et al**, (September 2005) *Is there a relationship between cell phone use and semen quality?*, *Arch Androl*. 2005 Sep-Oct;51(5):385-93
- Ferreira A et al**, (December 2006) *Ultra high frequency-electromagnetic field irradiation during pregnancy leads to an increase in erythrocytes micronuclei incidence in rat offspring*, *Life Sci* 2006 Dec 3;80(1):43-50

Friedman J et al, (August 2007) *Mechanism of a short-term ERK activation by electromagnetic fields at mobile phone frequency*, *Biochem J*. 2007 Aug 1;405(3):559-68

Fritzer G et al, (May 2007) *Effects of short- and long-term pulsed radiofrequency electromagnetic fields on night sleep and cognitive functions in healthy subject*, *Bioelectromagnetics*. 2007 May;28(4):316-25

George DF et al, (May 2008) *Non-thermal effects in the microwave induced unfolding of proteins observed by chaperone binding*, *Bioelectromagnetics*. 2008 May;29(4):324-30

Ghandi O, Kang G, (1996) *Effect of the head size on SAR for mobile telephones at 835 and 1900MHz*, *Bioelectromagnetics Society 23rd Annual Meeting*. St. Paul, Minnesota, USA, June 10-14, 2001, p. 52

Ghandi O, Kang G, (May 2002) *Some present problems and a proposed experimental phantom for SAR compliance testing of cellular telephones at 835 and 1900 MHz*, *Phys. Med. Biol.* 47 1501 18

Grajewski B et al, (October 2000) *Semen quality and hormone levels among radiofrequency heater operators*, *J Occup Environ Med*. 2000 Oct;42(10):993-1005

Haarala C et al, (May 2007) *Pulsed and continuous wave mobile phone exposure over left versus right hemisphere: Effects on human cognitive function*, *Bioelectromagnetics* 2007 May;28(4):289-95

Hardell L et al, (July 1999) *Use of cellular telephones and the risk for brain tumours: A case-control study*, *Int J Oncol*. 1999 Jul;15(1):113-6

Hardell L et al, (May 2000) *Case-control study on radiology work, medical x-ray investigations, and use of cellular telephones as risk factors for brain tumors*, *MedGenMed*. 2000 May 4;2(2):E2

Hardell L et al, (August 2002) *Cellular and cordless telephones and the risk for brain tumours*, *Eur J Cancer Prev*. 2002 Aug;11(4):377-86

Hardell L et al, (February 2003) *Further aspects on cellular and cordless telephones and brain tumours*, *Int J Oncol*. 2003 Feb;22(2):399-407

Hardell L et al, (March 2003) *Vestibular schwannoma, tinnitus and cellular telephones*, *Neuroepidemiology* 2003 Mar-Apr;22(2):124-9

Hardell L et al, (August 2004) *No association between the use of cellular or cordless telephones and salivary gland tumours*, *Occup Environ Med*. 2004 Aug;61(8):675-9

Hardell L et al, (2005) *Case-control study on cellular and cordless telephones and the risk for acoustic neuroma or meningioma in patients diagnosed 2000-2003*, *Neuroepidemiology*. 2005;25(3):120-8

Hardell L et al, (September 2005) *Use of cellular or cordless telephones and the risk for non-Hodgkin's lymphoma*, *Int Arch Occup Environ Health*. 2005 Sep;78(8):625-32

Hardell L et al, (February 2006) *Case-control study of the association between the use of cellular and cordless telephones and malignant brain tumors diagnosed during 2000-2003*, *Environ Res*. 2006 Feb;100(2):232-41

Hardell L et al, (October 2006) *Tumour risk associated with use of cellular telephones or cordless desktop telephones*, *World J Surg Oncol* 2006 Oct 11;4:74

- Hardell L et al**, (April 2007) *Use of cellular and cordless telephones and risk of testicular cancer*, Int J Androl. 2007 Apr;30(2):115-22
- Hardell L et al**, (September 2007) *Long-term use of cellular phones and brain tumours - increased risk associated with use for > 10 years*, Occup Environ Med. 2007 Sep;64(9):626-32
- Hepworth SJ et al**, (April 2006) *Mobile phone use and risk of glioma in adults: case-control study*, BMJ. 2006 Apr 15;332(7546):883-7
- Hocking B, Westerman R**, (September 2001) *Neurological abnormalities associated with CDMA exposure*, Occup Med (Lond). 2001 Sep;51(6):410-3
- Hocking B, Westerman R**, (October 2002) *Neurological changes induced by a mobile phone*, Occup Med (Lond). 2002 Oct;52(7):413-5
- Hocking B, Westerman R**, (March 2003) *Neurological effects of radiofrequency radiation*, Occup Med 2003 Mar;53(2):123-7
- Hours M et al**, (October 2007) *Cell Phones and Risk of brain and acoustic nerve tumours: the French INTERPHONE case-control study*, Rev Epidemiol Sante Publique. 2007 Oct;55(5):321-32
- Hoyto A et al**, (June 2007) *Ornithine decarboxylase activity is affected in primary astrocytes but not in secondary cell lines exposed to 872 MHz RF radiation*, Int J Radiat Biol. 2007 Jun;83(6):367-74
- Huber R et al**, (October 2000) *Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG*, Neuroreport. 2000 Oct 20;11(15):3321-5
- Huber R et al**, (December 2002) *Electromagnetic fields, such as those from mobile phones, alter regional cerebral blood flow and sleep and waking EEG*, J Sleep Res 2002 Dec;11(4):289-95
- Huber R et al**, (February 2005) *Exposure to pulse-modulated radio frequency electromagnetic fields affects regional cerebral blood flow*, Eur J Neurosci. 2005 Feb;21(4):1000-6
- Hung CS et al**, (June 2007) *Mobile phone 'talk-mode' signal delays EEG-determined sleep onset*, Neurosci Lett. 2007 Jun 21;421(1):82-6
- Inskip PD et al**, (January 2001) *Cellular-telephone use and brain tumors*, N Engl J Med. 2001 Jan 11;344(2):79-86
- Johansen C et al**, (February 2001) *Cellular telephones and cancer--a nationwide cohort study in Denmark*, J Natl Cancer Inst. 2001 Feb 7;93(3):203-7
- Johansen C et al**, (February 2002) *Mobile phones and malignant melanoma of the eye*, Br J Cancer. 2002 Feb 1;86(3):348-9
- Johansen C**, (2004) *Electromagnetic fields and health effects--epidemiologic studies of cancer, diseases of the central nervous system and arrhythmia-related heart disease*, Scand J Work Environ Health. 2004;30 Suppl 1:1-30
- Joubert V et al**, (January 2008) *Apoptosis is Induced by Radiofrequency Fields through the Caspase-Independent Mitochondrial Pathway in Cortical Neurons*, Radiat Res. 2008 Jan;169(1):38-45
- Kan P et al**, (January 2008) *Cellular phone use and brain tumor: a meta-analysis*, J Neurooncol. 2008 Jan;86(1):71-8

- Klaeboe L et al**, (April 2007) *Use of mobile phones in Norway and risk of intracranial tumours*, Eur J Cancer Prev. 2007 Apr;16(2):158-64
- Koivisto M et al**, (February 2000) *Effects of 902 MHz electromagnetic field emitted by cellular telephones on response times in humans*, Neuroreport. 2000 Feb 7;11(2):413-5
- Koivisto M et al**, (June 2000) *The effects of electromagnetic field emitted by GSM phones on working memory*, Neuroreport. 2000 Jun 5;11(8):1641-3
- Kramarenko AV, Tan U**, (July 2003) *Effects of high-frequency electromagnetic fields on human EEG: a brain mapping study*, Int J Neurosci. 2003 Jul;113(7):1007-19
- Krause CM et al**, (March 2000) *Effects of electromagnetic field emitted by cellular phones on the EEG during a memory task*, Neuroreport. 2000 Mar 20;11(4):761-4
- Krause CM et al**, (December 2000) *Effects of electromagnetic fields emitted by cellular phones on the electroencephalogram during a visual working memory task*, Int J Radiat Biol. 2000 Dec;76(12):1659-67
- Krause CM et al**, (June 2006) *Mobile phone effects on children's event-related oscillatory EEG during an auditory memory task*, Int J Radiat Biol 2006 Jun;82(6):443-50
- Krause CM et al**, (May 2007) *Effects of pulsed and continuous wave 902 MHz mobile phone exposure on brain oscillatory activity during cognitive processing*, Bioelectromagnetics 2007 May;28(4):296-308
- Lahkola A et al**, (May 2005) *Selection bias due to differential participation in a case-control study of mobile phone use and brain tumors*, Ann Epidemiol. 2005 May;15(5):321-5
- Lahkola A et al**, (April 2007) *Mobile phone use and risk of glioma in 5 North European countries*, Int J Cancer. 2007 Apr 15;120(8):1769-75
- Lahkola A et al**, (August 2008) *Meningioma and mobile phone use--a collaborative case-control study in five North European countries*, Int J Epidemiol. 2008 Aug 2. [Epub ahead of print]
- Lai H et al**, (May 1989) *Low-level microwave irradiation and central cholinergic systems*, Pharmacol Biochem Behav. 1989 May;33(1):131-8
- Lai H et al**, (1994) *Microwave irradiation affects radial-arm maze performance in the rat*, Bioelectromagnetics. 1994;15(2):95-104
- Lantow M et al**, (September 2006) *Comparative study of cell cycle kinetics and induction of apoptosis or necrosis after exposure of human mono mac 6 cells to radiofrequency radiation*, Radiat Res. 2006 Sep;166(3):539-43
- Leszczynski D et al**, (May 2002) *Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: molecular mechanism for cancer- and blood-brain barrier-related effects*, Differentiation. 2002 May;70(2-3):120-9
- Lonn S et al**, (January 2004) *Incidence trends of adult primary intracerebral tumors in four Nordic countries*, Int J Cancer. 2004 Jan 20;108(3):450-5
- Lonn S et al**, (November 2004) *Mobile phone use and the risk of acoustic neuroma*, Epidemiology. 2004 Nov;15(6):653-9

- Lonn S et al**, (March 2005) *Long-term mobile phone use and brain tumor risk*, Am J Epidemiol. 2005 Mar 15;161(6):526-35
- Lonn S et al**, (October 2006) *Mobile phone use and risk of parotid gland tumor*, Am J Epidemiol. 2006 Oct 1;164(7):637-43. Epub 2006 Jul 3
- Manti L et al**, (May 2008) *Effects of Modulated Microwave Radiation at Cellular Telephone Frequency (1.95 GHz) on X-Ray-Induced Chromosome Aberrations in Human Lymphocytes In Vitro*, Radiat Res. 2008 May;169(5):575-83
- Mazor R et al**, (January 2008) *Increased levels of numerical chromosome aberrations after in vitro exposure of human peripheral blood lymphocytes to radiofrequency electromagnetic fields for 72 hours*, Radiat Res. 2008 Jan;169(1):28-37
- Muscat JE et al**, (December 2000) *Handheld cellular telephone use and risk of brain cancer*, JAMA. 2000 Dec 20;284(23):3001-7
- Muscat JE et al**, (April 2002) *Handheld cellular telephones and risk of acoustic neuroma*, Neurology. 2002 Apr 23;58(8):1304-6
- Nikolova T et al**, (October 2005) *Electromagnetic fields affect transcript levels of apoptosis-related genes in embryonic stem cell-derived neural progenitor cells*, FASEB J. 2005 Oct;19(12):1686-8
- Nylund R, Leszczynski D**, (September 2006) *Mobile phone radiation causes changes in gene and protein expression in human endothelial cell lines and the response seems to be genome- and proteome-dependent*, Proteomics 2006 Sep;6(17):4769-80
- Oktay MF, Dasdag S**, (2006) *Effects of intensive and moderate cellular phone use on hearing function*, Electromagn Biol Med. 2006;25(1):13-21
- Oktem F et al**, (July 2005) *Oxidative damage in the kidney induced by 900-MHz-emitted mobile phone: protection by melatonin*, Arch Med Res. 2005 Jul-Aug;36(4):350-5
- Oral B et al**, (November 2006) *Endometrial apoptosis induced by a 900-MHz mobile phone: preventive effects of vitamins E and C*, Adv Ther. 2006 Nov-Dec;23(6):957-73
- Panagopoulos D et al**, (January 2007) *Cell death induced by GSM 900-MHz and DCS 1800-MHz mobile telephony radiation*, Mutat Res. 2007 Jan 10;626(1-2):69-78
- Papageorgiou CC et al**, (April 2006) *Acute mobile phone effects on pre-attentive operation*, Neurosci Lett. 2006 Apr 10-17;397(1-2):99-103
- Preece AW et al**, (2005) *Effect of 902 MHz mobile phone transmission on cognitive function in children*, Bioelectromagnetics Suppl 7 S138-43
- Rao VS et al**, (March 2008) *Nonthermal effects of radiofrequency-field exposure on calcium dynamics in stem cell-derived neuronal cells: elucidation of calcium pathways*, Radiat Res. 2008 Mar;169(3):319-29
- REFLEX Report**, (December 2004) *Risk Evaluation of Potential Environmental Hazards From Low Frequency Electromagnetic Field Exposure Using Sensitive in vitro Methods*, A project funded by the European Union under the programme "Quality of Life and Management of Living Resources"
- Rezk AY et al**, (February 2008) *Fetal and neonatal responses following maternal exposure to mobile phones*, Saudi Med J. 2008 Feb;29(2):218-23

- Sadetzki S et al**, (February 2008) *Cellular Phone Use and Risk of Benign and Malignant Parotid Gland Tumors A Nationwide Case-Control Study*, *Am J Epidemiol*. 2007 Dec 6 [Epub ahead of print]
- Salford L et al**, (June 2003) *Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones*, *Environ Health Perspect* 2003 Jun;111(7):881-3; discussion A408
- Samkange-Zeeb F et al**, (May 2004) *Validation of self-reported cellular phone use*, *J Expo Anal Environ Epidemiol*. 2004 May;14(3):245-8
- Schoemaker MJ et al**, (October 2005) *Mobile phone use and risk of acoustic neuroma: results of the Interphone case-control study in five North European countries*, *Br J Cancer*. 2005 Oct 3;93(7):842-8
- Schuz J et al**, (March 2006) *Cellular phones, cordless phones, and the risks of glioma and meningioma (Interphone Study Group, Germany)*, *Am J Epidemiol*. 2006 Mar 15;163(6):512-20
- Schuz J et al**, (December 2006) *Cellular telephone use and cancer risk: update of a nationwide Danish cohort*, *J Natl Cancer Inst*. 2006 Dec 6;98(23):1707-13
- Schwarz C et al**, (May 2008) *Radiofrequency electromagnetic fields (UMTS, 1,950 MHz) induce genotoxic effects in vitro in human fibroblasts but not in lymphocytes*, *Int Arch Occup Environ Health*. 2008 May;81(6):755-67
- Singh B, Bate LA**, (November 1996) *Responses of pulmonary intravascular macrophages to 915-MHz microwave radiation: ultrastructural and cytochemical study*, *Anat Rec*. 1996 Nov;246(3):343-55
- Stang A et al**, (January 2001) *The possible role of radiofrequency radiation in the development of uveal melanoma*, *Epidemiology*. 2001 Jan;12(1):7-12
- Takebayashi T et al**, (December 2006) *Mobile phone use and acoustic neuroma risk in Japan*, *Occup Environ Med*. 2006 Dec;63(12):802-7
- Takebayashi T et al**, (February 2008) *Mobile phone use, exposure to radiofrequency electromagnetic field, and brain tumour: a case-control study*, *Br J Cancer*. 2008 Feb 12;98(3):652-9
- Vrijheid M et al**, (April 2006) *Validation of short term recall of mobile phone use for the Interphone study*, *Occup Environ Med*. 2006 Apr;63(4):237-43
- Vrijheid M et al**, (May 2008) *Recall bias in the assessment of exposure to mobile phones*, *J Expo Sci Environ Epidemiol*. 2008 May 21
- Wiert J et al**, (July 2008) *Analysis of RF exposure in the head tissues of children and adults*, *Phys Med Biol*. 2008 Jul 7;53(13):3681-95
- Valbonesi P et al**, (March 2008) *Evaluation of HSP70 Expression and DNA Damage in Cells of a Human Trophoblast Cell Line Exposed to 1.8 GHz Amplitude-Modulated Radiofrequency Fields*, *Radiat Res*. 2008 Mar;169(3):270-9
- Velizarov S et al**, (February 1999) *The effects of radiofrequency fields on cell proliferation are non-thermal*, *Bioelectrochem Bioenerg*. 1999 Feb;48(1):177-80
- Wang Q et al**, (July 2004) *Effect of 900MHz electromagnetic fields on energy metabolism of cerebral cortical neurons in postnatal rat*, *Wei Sheng Yan Jiu*. 2004 Jul;33(4):428-9, 432

Wang Q et al, (March 2005) *Effect of 900Mhz electromagnetic fields on energy metabolism in postnatal rat cerebral cortical neurons*, Wei Sheng Yan Jiu. 2005 Mar;34(2):155-8

Wang Q et al, (September 2005) *Effect of 900 MHz electromagnetic fields on the expression of GABA receptor of cerebral cortical neurons in postnatal rats*, Wei Sheng Yan Jiu. 2005 Sep;34(5):546-8

Yan JG et al, (2008) *Upregulation of specific mRNA levels in rat brain after cell phone exposure*, Electromagn Biol Med. 2008;27(2):147-54

Yao K et al, (May 2008) *Effect of superposed electromagnetic noise on DNA damage of lens epithelial cells induced by microwave radiation*, Invest Ophthalmol Vis Sci. 2008 May;49(5):2009-15

Yao K et al, (May 2008) *Electromagnetic noise inhibits radiofrequency radiation-induced DNA damage and reactive oxygen species increase in human lens epithelial cells*, Mol Vis. 2008 May 19;14:964-9

ⁱ The DoH document, entitled 'Mobile phones and health', provides advice from the UK Chief Medical Officers who: "strongly advise that where children and young people do use mobile phones, they should be encouraged to:

- use mobile phones for essential purposes only
- keep all calls short – talking for long periods prolongs exposure and should be discouraged"

The advice is based on the findings of the Stewart Report, published in 2000, which was commissioned by government and led by a group of independent experts headed by Sir William Stewart. The report "...recommends 'a precautionary approach' to the use of mobile phones until more research findings become available...The expert group has therefore recommended that in line with a precautionary approach, the widespread use of mobile phones by children (under the age of 16) should be discouraged for non-essential calls."

ⁱⁱ Protective effects – a claim that the mobile phone protects the user from cancer

ⁱⁱⁱ Odds Ratio (OR) – where 1.0 = no risk, 2.0 = doubling of risk and 0.5 = halving of risk

^{iv} CI – Confidence interval. The band within which there is 95% confidence the risk applies

^v Electroencephalography (EEG) is the measurement of [electrical](#) activity produced by the [brain](#) as recorded from [electrodes](#) placed on the scalp

^{vi} Cerebral blood flow, or CBF, is the blood supply to the [brain](#) in a given time. In an adult, CBF is 750 mls/min or 15% of the cardiac output. On a weight basis, this is 50 to 54 millilitres/100grams/minute

^{vii} Double blind – where neither the participant nor the tester are aware whether the signal is real, to prevent bias

^{viii} Gene expression is the process by which inheritable information from a gene, such as the DNA sequence, is made into a functional gene product, such as protein or RNA